

Robust gamma oscillations in networks of inhibitory Hippocampal interneurons.

P.H.E. Tiesinga^{#,◇,*} and Jorge V. José[#]

[#] Center for Interdisciplinary Research on Complex Systems, and Department of Physics, Northeastern University, 360 Huntington Ave, Boston, Massachusetts 02115, USA

[◇] Sloan Center for Theoretical Neurobiology, Salk Institute,
10010 N. Torrey Pines Rd., La Jolla, CA 92037.

Recent experiments suggest that inhibitory networks of interneurons can synchronize the neuronal discharge in *in vitro* hippocampal slices. Subsequent theoretical work has shown that strong synchronization by mutual inhibition is only moderately robust against neuronal heterogeneities in the current drive, provided by activation of metabotropic glutamate receptors. *In vivo* neurons display greater variability in the interspike intervals due to the presence of synaptic noise. Noise and heterogeneity affect synchronization properties differently. In this paper we study using model simulations how robust synchronization can be in the presence of synaptic noise and neuronal heterogeneity. We find that with at least a minimum amount of noise stochastic weak synchronization (SWS) (i.e. when neurons spike within a short interval from each other, but not necessarily at each period) is produced that is much more robust than strong synchronization (i.e. when neurons spike each period). The statistics of the SWS population discharge are consistent with previous experimental data. We find robust SWS in the gamma frequency range (20-80 Hz) for a stronger synaptic coupling compared to previous models and for networks with 10 – 1000 neurons.

Keywords: interneuron, hippocampus, synchronization, noise, heterogeneity

*Corresponding author,
Tel: 619 453 4100 ext 1039
Fax: 619 455 7933
E-mail: tiesinga@salk.edu

I. INTRODUCTION

One of the important properties of the behavior of the nervous system, discovered early on [1], that has attracted significant amount of attention, is the synchronization of neuronal discharges. In recent years the advent of improved experimental techniques has provided vast amounts of new synchronization data. Concomitantly, there has been a resurgence in interest and controversy concerning the functional relevance of synchronization. It has been established that *in vivo* cortical neurons have noisy spike trains [2] (but see in contrast [3]), and that groups of neurons discharge coherently as found in population recordings (such as EEGs, or by arrays of extracellular electrodes, for a review see [4]). These two facts have sparked major controversies. Firstly, does noise (or precise timing) in neuronal spike trains contain information [5,6], or is information merely due to noisy processing of an average firing rate [7–9]? Secondly, is synchronization functionally (or even statistically) significant [10,11], or just an epiphenomenon [9]? In this paper we focus on two different aspects of synchronization that have received little attention so far. Can realistic neuronal networks synchronize under the biological conditions of variable intrinsic neuronal properties, and the noise-induced neuronal unreliability? What kind of synchronization can be obtained, and what are its pertinent statistical properties? It is necessary to resolve these two questions to properly formulate the issues to be studied in experiment, and to analyze different ways of probing the experimental data. Here we focus our attention on the extensively-studied synchronous gamma oscillations in hippocampus [12–17]. Theoretical and computational work has shown that mutual inhibition is capable of synchronizing neuronal networks [18,19]. Subsequent *in vitro* experiments have convincingly established the role of GABA-ergic hippocampal interneurons in gamma oscillations [14,12]. Wang & Buzsáki [20] studied the effect of current heterogeneity and partial connectivity on the synchronization of the hippocampal network. They only found strong synchronization in the gamma frequency range when the current heterogeneities were small [20,21]. In strong synchronization all neurons in a local circuit spike within a short interval of each other. This suggests that strong synchronization can only be obtained when the intrinsic properties of the neurons are not too different. According to Ref. [20] this would mean a less than 10% difference in current drive, or average firing rate. It has been hard to pinpoint the amount of variability in intrinsic properties in the *in vitro* and *in vivo* preparations of different brain areas. It is however not unreasonable to assume the presence of more than 10% variability in these preparations. Strong synchronization is also not robust against noise [22]. It would therefore seem unlikely for strong synchronization to be present in hippocampus under physiological conditions. Indeed, here we show that stochastic weak synchronization (SWS) is more prevalent in parameter space, and is also robust against neuronal heterogeneities and synaptic noise. We conjecture that as a consequence it is much more likely to occur in neuronal systems.

In SWS, neurons spike within a short interval from each other, but not necessarily at each period [23,24]. The synchronization is called stochastic, because the particular cycle in which the neurons fire is random. This makes the properties of this state different from the well known cluster states studied by previous authors [25–28]. There each neuron always fires at the same cycle with the same cluster. Both strong and stochastic weak synchronization yield periodic population oscillations. The difference can then only be ascertained using multi-unit recordings.

We use cross correlation analysis to show that noise and heterogeneity affect the synchronization properties of our network in very different ways. Large enough noise and heterogeneity will, however, stop strongly synchronized oscillations. We demonstrate that neither adding a periodic drive nor increasing synaptic coupling can significantly increase robustness of strong synchronization. Finally we determine for what parameters robust self-induced 40Hz synchronous oscillations can be obtained.

II. METHODS

A. Single neuron model

Our aim here is to establish physiological criteria for robust synchronization in the gamma frequency range. The use of a biophysically realistic model is therefore of pivotal importance. At the same time it is also important to balance the amount of complexity versus practical simplicity [29]. We have therefore not attempted to use the latest available data to construct a detailed multi-compartmental model. The computer requirements to sample the full relevant parameter space, and perform our type of analysis, would be extremely demanding even using very fast computers. It has been shown, nonetheless, that one and two compartmental models can accurately generate spike trains of the right shape and frequency [30–32]. Multi-compartmental models may be necessary to assess the synaptic integration of inputs located on different parts of the dendritic tree. This is currently an intensely studied area in electrophysiology [33–35]. Here we study a model previously introduced by others [20]. The model has been shown to reproduce the salient features of the dynamics of hippocampal interneurons. The neurons are modeled as a single compartment with Hodgkin-Huxley type sodium and potassium channels. In this work all the neurons are connected to all others and themselves (ALL to ALL connectivity) via inhibitory GABA_A-synapses. The equation for the membrane potential of a neuron is (the index i of the neuron is omitted)

$$C_m \frac{dV}{dt} = -I_{Na} - I_K - I_L - I_{syn} + I + C_m \xi. \quad (1)$$

Here we use: the leak current $I_L = g_L(V - E_L)$, the sodium current $I_{Na} = g_{Na}m_\infty^3 h(V - E_{Na})$, the potassium current: $I_K = g_K n^4(V - E_K)$, and the synaptic current: $I_{syn} = g_{syn}s(V - E_{syn})$. The Gaussian noise is denoted as ξ (see below), and I is the tonic drive. The channel kinetics are given

in terms of m , n , and h . They satisfy the following first order kinetics:

$$\frac{dx}{dt} = \phi(\alpha_x(1 - x) - \beta_x x). \quad (2)$$

Here x labels the different kinetic variables m , n , and h , and $\phi = 5$ is a dimensionless time-scale that can be used to tune the temperature-dependent speed with which the channels open or close. The rate constants are [20],

$$\begin{aligned} \alpha_m &= \frac{-0.1(V + 35)}{\exp(-0.1(V + 35)) - 1}, \\ \beta_m &= 4 \exp(-(V + 60)/18), \\ \alpha_h &= 0.07 \exp(-(V + 58)/20), \\ \beta_h &= \frac{1}{\exp(-0.1(V + 28)) + 1}, \\ \alpha_n &= \frac{-0.01(V + 34)}{\exp(-0.1(V + 34)) - 1}, \\ \beta_n &= 0.125 \exp(-(V + 44)/80). \end{aligned}$$

We make the approximation that m follows the asymptotic value $m_\infty(V(t)) = \alpha_m/(\alpha_m + \beta_m)$ instantaneously. The synaptic gating variable s obeys the following equation [36,37,20]:

$$\frac{ds}{dt} = \alpha F(V_p)(1 - s) - \beta s, \quad (3)$$

with $\alpha = 12 \text{ ms}^{-1}$, $\beta = 1/\tau_{syn}$, $F(V_p) = 1/(\exp(-V_p/2) + 1)$, and V_p is the presynaptic potential. The function $F(V_p)$ is chosen such that when the presynaptic neuron fires, $V_p > 0$, the synaptic channel opens. The decay time of the postsynaptic hyperpolarization is chosen as $\tau_{syn} = 1/\beta = 10 \text{ ms}$ (or 20 ms in some instances). We use a reversal potential of $E_{syn} = -75 \text{ mV}$ for the inhibitory (GABA_A) synapses [38]. The standard set of values for the conductances used in this work is $g_{Na} = 35$, $g_K = 9$, $g_L = 0.1$, and $g_{syn} = 0.1$ (in mS/cm^2), and we have taken $E_{Na} = 55 \text{ mV}$, $E_K = -90 \text{ mV}$, and $E_L = -65 \text{ mV}$. The membrane capacitance is $C_m = 1 \mu\text{F}/\text{cm}^2$. Unless stated otherwise we will use the standard set of parameters listed above. When no current value is specified we use $I = 1 \mu\text{A}/\text{cm}^2$. The network will then spike at approximately 39 Hz.

We chose the initial values for the membrane potential at the start of the simulations uniformly random between -70 and -50 mV . The kinetic variables m , n , h , and s are set to their asymptotic stationary values corresponding to that starting value of the membrane potential.

The resulting equations with noise are integrated using an adapted second order Runge-Kutta method [39], with time step $dt = 0.01 \text{ ms}$. The accuracy of this integration method was checked for the dynamical equations without noise ($D = 0$) by varying dt and comparing the result to the one obtained with the standard 4th order Runge-Kutta method [40] with a time-step dt of 0.05 ms .

We normalize all quantities by the surface area of the neuron. This leads to the following system of units: the membrane potential V in mV , time t in ms , firing rate f in Hz ,

membrane capacitance C_m in $\mu\text{F}/\text{cm}^2$, conductance g_x in mS/cm^2 , voltage noise ξ in mV/ms , strength of neuroelectric noise D in mV^2/ms , the rate constants α_x and β_x in ms^{-1} , and the current I in $\mu\text{A}/\text{cm}^2$. The kinetic variables m , n , h , s , and the time-scale ϕ are dimensionless. Results in our paper are expressed in this system of units.

B. Heterogeneity and synaptic noise

We have included heterogeneity in the applied current. For each run we draw the applied current for each neuron from a uniform distribution. The average of the current distribution is I and the variance is σ_I^2 . The current heterogeneity represents the variation in the intrinsic properties of the neurons in the hippocampus. Experimental measurements of quantities like the input resistance R_{in} , the membrane time-scale, the spontaneous spiking rate, the shape of the somatic action potential (amplitude, width, rise and fall time), and the afterhyperpolarization, show considerable variance [41–43]. It is hard to determine how much of the variance is due to measurement errors, and how much is actually due to intrinsic neuronal variability. Here we assume the main effect of the variability is to change the intrinsic frequency of the neurons (which can be varied using the current drive in our model). Another source of heterogeneity in *in vitro* experiments is the glutamate pressure ejection method [14]. It can lead to an inhomogeneous activation of metabotropic glutamate receptors, and thus to a variable current. In this paper we will consider σ_I as a free parameter.

At least three sources of noise can be identified [44]: random inhibitory postsynaptic potentials (IPSP) and excitatory postsynaptic potentials (EPSP), stochasticity of the synaptic transmission, and the stochasticity of the channel dynamics. Here we assume that the variability in the neuronal discharge is mainly due to synaptic noise [45]. We have compared the effects of Poisson distributed spike trains of EPSPs and IPSPs to that of a Gaussian noise current on interspike interval (ISI) variability. Poisson and Gaussian noises, do not yield identical results. The statistics obtained from both models, however, are similar in the parameter regime studied [46]. For the purpose of our studies we consider that Poisson and Gaussian distributions are two alternate ways of producing noisy spike trains with particular statistics. Therefore, the synaptic noise is only implemented as a Gaussian distributed, white noise current in neuron i , with $\langle \xi_i(t) \rangle = 0$, and $\langle \xi_i(t) \xi_j(t') \rangle = 2D\delta(t - t')\delta_{ij}$. The noise currents in different neurons are assumed independent.

C. Measured quantities

From our simulations we obtain the time trace for the membrane potential $V_i(t)$ of each neuron. We determine the spike-trace X_i from V_i as follows: $X_i(t) = 1$ when $V_i(t)$ crosses 0 mV (i.e. $V(t^-) \leq 0 < V(t^+)$), and it is zero elsewhere. From X_i we obtain $X(t) = \sum_i X_i(t)$. X is proportional to

the instantaneous firing rate of the network. We also calculate the correlations function κ [20]:

$$\kappa \equiv \sum_{i \neq j} \frac{\langle \hat{X}_i(n\tau) \hat{X}_j(n\tau) \rangle}{\sqrt{\langle \hat{X}_i(n\tau) \rangle \langle \hat{X}_j(n\tau) \rangle}}. \quad (4)$$

This function measures the amount of strong synchronization, and depends on the bin size τ of the time discretization

$$\hat{X}_i(n\tau) = \int_{(n-1)\tau}^{n\tau} ds X_i(s). \quad (5)$$

We use $\tau = 200 \text{ dt} = 2 \text{ ms}$ for oscillations in the gamma-frequency range, or $T/10$ for periodic drives with period T .

We also evaluate other measures that yield further detailed quantitative characterization of the network behavior. We calculate the time autocorrelation function:

$$g_x(t) = \frac{\langle x(t)x(0) \rangle - \langle x \rangle^2}{(\langle x^2 \rangle - \langle x \rangle^2)}, \quad (6)$$

and the cross correlation function:

$$g_{xy}(t) = \frac{\langle x(t)y(0) \rangle - \langle x(t) \rangle \langle y(0) \rangle}{\sqrt{\langle x(t)^2 \rangle \langle y(0)^2 \rangle}}. \quad (7)$$

Here x and y can be any of the variables X_i , V_i , and X , and $\langle \cdot \rangle$ is a shorthand notation for the time-average.

We also consider the more conventional interspike interval histogram (ISIH) [47], averaged over all network neurons. From the ISIH one can obtain two statistics: the average ISI, τ_{ISI} , and the standard deviation of the ISI, σ_{ISI} . The ratio σ_{ISI}/τ_{ISI} is known as the coefficient of variation (CV). The average firing rate is $f = 1/\tau_{ISI}$, and the population standard deviation of f is σ_f .

$$\sigma_f = \sum_j f_j^2 - f^2, \quad (8)$$

where $f_j = 1/\tau_{ISI}^j$ is the average firing rate of the j th neuron. In addition we plot rastergrams, with the action potential of each neuron plotted as a filled circle, with the y-coordinate given by the neuron index and the x-coordinate by the spiking time.

To analyze the stochastic weak synchronization network dynamics we need to apply a different method. The population period τ_n is different from the population averaged ISI, and to estimate it we proceed as follows. First we determine the firing rate $\hat{X}(t)$ as before with 1 ms bins. In the stochastic weak synchronization state $\hat{X}(t)$ will consist of a number of approximately equidistant peaks of finite width (see Fig. 11f). We use the position of the first maximum of the Fourier transform at nonzero frequency as an estimate T for the period τ_n . We calculated the weight $\langle \hat{X}(t) \rangle$, the average position $t_c^i = \langle t \hat{X}(t) \rangle$, and the width $\sigma_c^i = \sqrt{\langle t^2 \hat{X}(t) \rangle - (t_c^i)^2}$ of the i th peak. The time average is taken over a range $[-0.35T, 0.35T]$ about the estimated position $t_c^{i-1} + T$ of

the peak. We calculated the number of spikes that fall outside this region. If the average number of missed spikes is more than one per cycle we reject the cluster state. The cycle length (time between two consecutive cluster firings) is defined as $\tau_c^i = t_c^i - t_c^{i-1}$. We determine the average cluster size $N_c = \langle N_c^i \rangle$, its $CV(N_c) = \sqrt{\langle (N_c^i)^2 \rangle - N_c^2}/N_c$, the average cycle length $\tau_n = \langle \tau_c^i \rangle$, its $CV(\tau_n) = \sqrt{\langle (\tau_c^i)^2 \rangle - \tau_n^2}/\tau_n$, and the average width $\sigma_c = \langle \sigma_c^i \rangle$. Here the average $\langle \cdot \rangle$ is given by the sum over all cycles in the run (after discarding a transient). We characterize the strength of the synchronization using a modified κ_W and CV_W . In the SWS state the ISIH has multiple peaks. The CV of the ISI receives contributions from the variance within each peak, but also of the variance between the multiple peaks. We are only interested in the former, and the conventional CV is thus an overestimate. Instead we use $CV_W = \sigma_c/\tau_n$ which is related to the average width of one peak in the ISIH. The coherence κ measures the number of coincident spikes between two spike trains. Consider two neurons that do not spike at each cycle, but when they both do, the spikes are coincident (that is in the same bin). If the probability of spiking in a cycle is $p = N_c/N$, and both neurons fire statistically independent, we obtain $\kappa = p$. These neurons can be considered synchronous and we want $\kappa_W = 1$. We therefore normalize κ by p .

There is a subtlety in the calculation of the average firing rate. In the deterministic noiseless case one ISI is enough to determine the average value (after discarding the transient). [Note that counting the number of spikes in a fixed interval is not an efficient way to determine the exact firing rate.] In the presence of noise, however, you need at least 10 ISIs to accurately determine the average. In networks with large current heterogeneities there are neurons with high and very low firing rates (Fig. 11). The average ISI for the low firing rate is less accurate than for the high firing rate neurons in the network. However, it carries equal weight in the conventional average $\tau_{ISI} = \sum_j \tau_{ISI}^j$. We have therefore used a weighted average $\tau_{ISI} = \sum_j n_j \tau_{ISI}^j / \sum_j n_j$ (here n_j is the number of intervals over which τ_{ISI}^j is calculated), and the approximate identity $N_c/\tau_n \approx N_s/\tau_{ISI}$ can be used as a check. N_s is the number of active neurons, defined as the neurons that have more than two ISIs after the transient.

III. RESULTS

A. Non robustness of strong synchronization

In this section we describe the results of our simulations for a network of $N = 100$ interneurons, connected all to all, with either synaptic noise (SN), or current heterogeneities (CH). In Fig. 1 we plot coherence parameter κ (defined in Eq. (4)) versus the strength of the synaptic noise D , and versus the standard deviation of the current heterogeneities σ_I . We find that strong synchronization is lost for approximately $D > 0.10 \text{ mV}^2/\text{ms}$ and $\sigma_I > 0.1 \text{ } \mu\text{A}/\text{cm}^2$ (with the standard set of parameters listed in Methods). The mechanism by

which strong synchronization is lost, however, is different in the CH case compared to the mechanism with SN. This difference shows up only if we studies the whole state of the network using cross correlation functions, instead of the average quantities shown in Fig. 1. Next we compare these two mechanisms. Wang&Buzsaki (WB) [20] have already analyzed the case with current heterogeneity. We have reproduced part of their work, and we will refer to their corresponding figures. In both CH and SN cases the neuronal firing rate decreases when the network desynchronizes. We have plotted the time-trace of the synaptic drive $s(t)$ in Figs. 1c and d. The phasic part decreases, and the tonic part of $s(t)$ increases with increasing D and σ_I . The increased tonic part is responsible for the lower average firing rate. The firing rate of the CH neurons saturates (when averaged over enough realizations of the current heterogeneities¹), whereas for the SN it increases steadily as a function of D for large values of D . This is because the single neuron firing rate increases with D [46], but tonic inhibition saturates to its highest value in the asynchronous network. The dispersion σ_f (see Methods) with CH is larger than the one in SN (not shown). In SN all the neurons have identical intrinsic properties, and the expectation value for the average frequency of each neuron is the same. The dispersion σ_f in this case represents the fluctuations in the average ISI due to the finite averaging time. With CH the neurons have different intrinsic frequencies, and the dispersion σ_f increases with σ_I (and does not go to zero after a long averaging time, see WB Fig. 5B). In Fig 2I and II we compare the correlation functions for the SN and CH case, respectively. In (a) we have the strongly synchronous network, in (c) the asynchronous network, and in (b) a transition state. The difference between SN and CH becomes clear when one considers the cross correlation functions. With CH the number of pairs that are phase locked drops gradually (see WB Fig. 8E). The pairs that are phase locked, are tightly phase locked (Fig. 2Ib1,4,5,6), and there is no dispersion in the cross correlations, only a relative phase. Even in the asynchronous state the autocorrelation function g_{X_i} for a single neuron is sharp, i.e. the neuron fires regularly with a fixed frequency (Fig. 2Ic1 and c13). The population average of g_{X_i} , however, is disordered (Fig. 2Ic12), since each of the neurons has a different firing rate. In the SN case there is already dispersion due to the noise-induced jitter in the spike time, in the autocorrelation Fig. 2IIa11, and in the cross correlations (Fig. 2IIa2-a10). The dispersion increases gradually with D . The difference between the CH and SN cases is also evident in the distribution of κ values for each pair in the network (Fig. 3, WB Fig. 8E). For SN there is a well defined peak, with the average shifting to lower values as D increases, (Fig. 3a-c), whereas for CH there is a broad

distribution for small σ_I (Fig. 3d), a peak at low values of κ combined with a broad distribution for moderate values of σ_I (Fig. 3e). For higher values of σ_I the network is in an asynchronous regime, and only the peak for low κ values is present (Fig. 3f). We have compared the ISIH for a network neuron to the ISIH of an isolated neuron (Figs. 4c-e), and also the values of τ_{ISI} and σ_{ISI} (Figs. 4a and b). The CV of the network neuron is higher than the CV of an isolated neuron which in turn is higher than the CV of an isolated neuron with aut synaptic feedback. The inhibitory coupling in the network increases the effect of the noise compared to uncoupled neurons: the jitter in the spike times reduces the phasic component of $s(t)$ (Fig. 1c). This effect does not take place in a neuron with aut synaptic feedback: the size of the phasic component does not decrease with D , only the timing deteriorates.

B. Weak enhancement of robustness due to resonance effect

In this section we drive the tonically active neurons with an external periodic drive. This drive may represent the effects of putative pacemaker neurons, similar to the ones that were recently found in the striate cortex of cats [48]. There is no compelling evidence for having the 40Hz pacemaker neurons projecting to hippocampal interneurons. Here we model the pacemaker as an excitatory synapse driven by a periodic pulse train, and investigate its effects on the robustness of synchronization. When a single neuron is driven by a periodic drive it will entrain, or phase lock, when the drive frequency is close to the natural frequency (or in some cases close to a rational fraction) [49]. Here we vary the natural frequency via the current. In Fig. 5c we show the coefficient of variation (CV) of an isolated neuron. When the neuron is entrained the CV drops to zero. The ISIH then consists of a single peak. The entrainment occurs for a range of current values, $I = 0.87-1.0$. Since the firing rate is constant, the $f - I$ has a flat step (not shown). Outside the range of entrainment the ISIH has more structure (Fig. 5d and f). When there is noise present in the neuron, the CV will increase. For weak noise the CV in the entrainment regime will still be lower compared to the CV when the neuron is not entrained. For the network the CV is also lower in the entrainment regime (Fig. 5b), though the CV increases faster with the noise strength compared to the CV of the isolated neuron. The synchronization of the network, measured by κ (Fig. 5c), is significantly enhanced in the region of entrainment (for $D = 0.004-0.04$). The enhanced synchronization disappears for higher values $D > 0.2$.

C. Effect of synaptic coupling strength on robustness of strong synchronization

We have also studied the effect of varying the synaptic coupling strength g_{syn} . For large enough D the network will be asynchronous. We find that the network frequency in that case decreases with increasing values of g_{syn} (Fig. 6). For an

¹This assumes

$$\int_{I_{av}-\sigma_I\sqrt{12}}^{I_{av}+\sigma_I\sqrt{12}} f(I)dI \approx f(I_{av})$$

asynchronous network the synaptic drive has a constant tonic hyperpolarizing conductance, decreasing the firing rate. The stronger the coupling the larger the decrease. The synchronization measured by the parameter κ displays a different behavior. In Fig. 6c we plot the κ versus g_{syn} for one specific value $D = 0.02$ (and I chosen such that the firing rate is approximately 39 Hz). It is interesting to note that stronger coupling does not necessarily mean a higher value of κ . The coherence κ has a local maximum for $g_{syn} = 0.1$, for higher values of g_{syn} , κ decreases (see WB Fig. 12B). For $g_{syn} > 0.3$, κ starts increasing again. We have studied the underlying dynamics of this non monotonous behavior. In Fig. 7c-f we plot the ISIH for different values of g_{syn} . For $g_{syn} \geq 0.2$ one finds more than one peak. In the rastergrams (Figs. 7a and (b)), one can see that the dynamics corresponds to a population that has a well defined frequency, but individual neurons sometimes miss, or skip, a period. Despite this small asynchrony when the neuron fires, it does so in synchrony with the others. As a consequence the rastergrams looks much more ordered compared to the one for $g_{syn} = 0.1$ at the same noise strength $D = 0.002$.

D. Larger g_{syn} leads to robust stochastic weak synchronization

In this subsection we discuss the robust 40 Hz rhythms found for higher g_{syn} values. We have doubled the synaptic decay constant to $\tau_{syn} = 20$. Here we will use the modified κ_W and CV_W as mentioned in the Methods section.

In Fig. 8 we vary g_{syn} from 0.05 to 2.5 with a spacing of 0.05. The neuron number is kept equal to $N = 100$, and we use $I = 2.0$, and $\sigma_I = 0$. For $D = 0.0$ and $\sigma_I = 0$ the network is in a strongly synchronized state, with the network frequency f_n the same as the single neuron firing rate f . The frequency is exactly the same as the one for a single neuron with autosynaptic feedback, as one would expect. This coherent state can be arrived at from many different random initial conditions.

For weak noise, $D = 0.008$, the network stays in a strongly synchronized state for $g_{syn} < 0.25$. For higher g_{syn} skipping starts to occur, the fractional cluster size decreases from values close to one to values below one-half at $g_{syn} = 1.2$. At that point the network is in a real (albeit stochastic) cluster state, on average the neuron only fires once every two cycles. We will refer to all states for which certain active neurons do not fire at each cycle as a stochastic weak synchronized (SWS) network. The network frequency, f_n , and the single neuron firing rate f both decrease with increasing g_{syn} . When the network settles in the SWS state f starts to differ considerably from its value at the $D = 0$ state. The strength of synchronization increases with g_{syn} , that is CV_W decreases and κ_W increases. For values $g_{syn} > 2.0$, CV_W and κ_W slowly saturate.

For stronger noise $D = 0.04$ and $D = 0.20$ the network is asynchronous for low values of g_{syn} . We have therefore excluded these points based on the criterium discussed in the Methods section. The network frequency starts out at a higher

value, and the neurons fire at a lower rate compared to the $D = 0.008$ case. The strength of synchronization, κ_W and CV_W , is reduced compared to the one for $D = 0.008$, but still increases with g_{syn} . Note that all the neurons in the network still have a nonzero firing rate. It is thus possible to obtain weakly synchronized oscillations in a network consisting of 100 neurons in the frequency range between 20 and 40 Hz.

We find that noise is necessary to obtain SWS. We have studied SWS in the presence of weak current heterogeneities, say for $\sigma_I = 0.02$. Without noise ($D = 0$) the network is in an strongly synchronized state, and κ displays a maximum as a function of g_{syn} (WB Fig12B). One also clearly notices the effect of suppression [21]: for larger g_{syn} the inhibition of faster spiking neurons stops the firing of neurons driven by a smaller current. As a result the total number N_s of active neurons gradually drops (Fig. 9f). For a small amount of noise, $D = 0.008$, the situation changes dramatically. A SWS state is obtained, and all neurons remain active ($N_s/N = 1$), while κ saturates for $g_{syn} > 0.5$, and the value of κ for $g_{syn} > 0.8$ is even higher than without noise. Thus noise may actually improve the coincidence. Of course noise does increase the width, CV_W , of the peaks in the instantaneous firing rate. The single neuron firing rate decreases significantly compared to that for $D = 0$, whereas the network frequency is only weakly affected.

We have performed numerical simulations for system sizes $N = 10, 20, 50, 100, 200$, and 1000. We have used the following parameters values: $I = 5.0$, $D = 0.2$, $g_{syn} = 1$, $\tau_{syn} = 20$, and $\sigma_I = 0.1$. The network frequency increases with system size, whereas the firing rate stays approximately constant with a dip around $N = 50$. The measures for coherence, κ_W and CV_W , are also only weakly dependent on system size.

The cycle-to-cycle fluctuations in cluster size vary approximately as $\sqrt{N}$ (Fig. 10g). The strength of the inhibition is determined by the number of neurons that fired in the previous cluster, and in turn it determines at what time the first neurons become disinhibited. One therefore expects cluster size fluctuations and cycle length fluctuations to be intimately related. Indeed, the standard deviation of cycle length varies as $1/\sqrt{N}$ with N the number of neurons (Fig. 10h). This means that larger networks are better at generating a precise cycle length, whereas size does not matter as much for the coincidence of spikes measured by κ_W and CV_W .

In each simulation we randomly draw a set of driving currents I_j for each neuron j from a uniform probability distribution. The results one obtains may critically depend on the particular realization of driving currents. One expects that for larger systems this is less of a problem. The population distribution of I is more likely to approach the original ensemble distribution of currents for a given neuron. Here we have studied the range of values for the measured quantities (f_n , f , and so on) for ten different realizations. We find that for most quantities (for these parameter values) the range of values decreases with N , and for $N \geq 500$ one realization will give a result close to the expectation value.

We now vary σ_I and D for the following fixed parameter set $N = 1000$, $g_{syn} = 2$, $\tau_{syn} = 20$, and $I = 3.5$. For

$D = 0$ and $\sigma_I = 0$ the network is strongly synchronized at 20 Hz. The instantaneous firing rate consists of a sequence of regularly spaced delta functions (Fig. 11e), the ISIH has a single delta peak at 50 ms (Fig. 11b), and all neurons spike at the same frequency (Fig. 11a). Increasing D increases the network frequency, but decreases the single neuron firing rate (Fig. 12). The population activity is still periodic (Fig. 11f), but the peaks have a finite width (as well as the ISIH), and the ISIH becomes multimodal. This process continues with ISIH spreading out more and more, with the CV_W increasing, and κ_W decreasing.

As mentioned before we need some noise to generate an SWS state. Here we use $D = 0.2$, while at the same time varying σ_I . For finite σ_I there is still a coherent population activity (Fig. 11g), despite the fact that neurons have different firing rates (Fig. 11c). Increasing σ_I will reduce coherence, κ_W decreases and CV_W increases. At the same time both f_n and f increase (Fig. 12d). This is different from the effect of increasing D . Higher σ_I leads to suppression, with fast spiking neurons preventing slower ones from firing, and as a result part of the inhibition disappears, while further increasing the firing rate and its average (calculated from the active neurons). On the other hand noise increases the tonic inhibition for each neuron, and thus leads to a reduced firing rate. Also the progression of the asynchronous state is different. The first peak in the ISIH becomes broader, and the higher order ones have a reduced prominence (Fig. 11d). For increasing D the peaks just wash out.

E. Comparison to externally induced stochastic weak synchronization

We find that robustness can be enhanced by getting a network in a SWS state. The single neuron discharge in an SWS state is very similar to that obtained in Stochastic Resonance [50] using a subthreshold *external* drive. In this subsection we compare the previous self-induced SWS state to the one induced by an external drive. We drive the system by a sinusoidal current of amplitude 1.2 (admittedly large). The average value of the driving current is zero. For weak noise the spikes of a single neuron are spaced many cycles apart (Fig. 13a). With increasing noise the ISIH starts to look more like the ones in Fig. 11b. The population activity also is periodic (Fig. 13e). The distribution of τ_{ISI} in Fig. 13a looks similar to the one in Fig. 11a. For high values of D the neuron can spike more than once during one cycle, as a result the number of coincident spikes is reduced. The addition of current heterogeneity does not seem to affect the network behavior (as long as all the neurons are still below threshold). The different ISIH in Fig. 13d look very similar. However the corresponding range of τ_{ISI} values in Fig. 11c does increase with σ_I .

IV. DISCUSSION

Previous authors have recognized that strong synchronization is only moderately robust against neuronal heterogeneity [20,21]. We have previously shown that the same holds including synaptic noise [22]. The basic premise of synchronization by mutual inhibition is almost trivial, since the network consists of intrinsically periodically spiking neurons. Their output produces a periodic synaptic drive, which in turn is fed back into the network. Inhibition thus allows a phase lock at zero phase with this drive. Heterogeneity and noise reduces the phasic, and increases the tonic part of the synaptic drive, leading to a reduction in synchronization, and eventually leading to an asynchronous state (Fig. 1c,d). The synchronization behavior of networks of physiological realistic neurons, however, is by no means fully understood. In this work we showed that the loss of synchronization proceeds via **different** mechanisms in the presence of synaptic noise compared to the presence of current heterogeneity. This is evident from the cross correlations shown in Figs. 2 and 3. We also found that the noise-induced precision loss in the uncoupled neuron is exacerbated by the inhibitory coupling. All of these could seem obvious based on previous work on heterogeneity [20,21]. However, its consequences for real life biological networks had not been fully appreciated. Our results, combined with previous results, show that there is a problem with strong synchronization by mutual inhibition, since it is unlikely to occur in *in vitro* or *in vivo* systems. [There are exceptions such as for example the pacemaker nucleus in electric fish [51], where the neurons are coupled via gap junctions.] The aim of this paper was to treat this problem: how can one obtain robust synchronization in the presence of synaptic noise and neuronal heterogeneity? Our results are twofold. First, methods to increase the robustness of strong synchronization have been ineffective. Second, we showed that robust stochastic weak synchronization (SWS) can be obtained for biophysically realistic parameter values. SWS is consistent with previous experimental data. In what follows we discuss these two important results in more detail.

We believe that strong synchronization is not robust enough. To make sure that we do not prematurely discard strong synchronization by mutual inhibition we have to make an effort to increase robustness. In this paper we discussed two simple methods to increase robustness of strong synchronization. One method was to increase the synaptic coupling g_{syn} . Inhibition is responsible for synchronization. It is then quite natural to expect that increasing the strength of inhibition increases robustness. The fact that this does not happen is surprising. For current heterogeneity this is in part due to suppression [21,20]. We have studied this effect for synaptic noise in more detail. We found that neurons skip periods for higher values of g_{syn} (see Fig. 7). In other words the strongly synchronized state becomes unstable, and a weakly synchronized state emerges. This weakly synchronized state looked more coherent (Fig. 7a,b), and provided the impetus to further study the robustness of the SWS states.

We also added a periodic drive to the neuron. Recent exper-

imental work shows that the CV of neurons on an entrainment step is reduced compared to the CV outside the step [49]. A clear physiological correlate in hippocampus of this drive is lacking at present. Since we try to reject our conjecture this lack of physiological realism is not a problem. We found a moderate increase in robustness (see Fig. 5). The periodic drive is not as effective as one would have intuited, however. In fact the inhibitory connections reduce the increase in robustness compared to the increase in the single uncoupled neuron (see Fig. 5). If we add a subthreshold periodic drive with noise to a quiescent neuron we obtain weak synchronization. This is known as Stochastic Resonance in excitable systems [50]. To summarize: our attempts to significantly increase robustness of strong synchronization failed. Instead we found weak synchronization. Thus it is easier to find weak synchronization in parameter space than it is to find strong synchronization.

If one accepts the fact, however, that strong synchronization is not robust against noise and heterogeneity, and that periodic population oscillations are found in experiments, then one has to carefully consider the possible relevance of weak synchronization. Weak synchronization as well as strong synchronization lead to a periodic population discharge, and specifically to an inhibitory synaptic drive indistinguishable from the one found in pyramidal neurons in [14]. Moreover, the clusters that form in stochastic weak synchronization bear a resemblance to the neuronal assemblies found in some experiments [52], and that are thought to play a role in putative binding [53]. The question then is, is SWS more robust, and can it be found for the gamma frequency range for biophysically realistic parameters? What is needed is a higher total synaptic conductance, and noise. The necessary amount of noise is very small, $D > 0.004$ is sufficient. The noise prevents the occurrence of suppression (Fig. 9). In suppression the faster neurons prevent the slower ones from firing. This reduces the inhibition of the faster ones, and allows them to fire at different frequencies and at random relative phases. Suppression is thus detrimental to synchronization.

We obtained SWS for different system sizes (we studied networks from 10 to 1000 neurons). The coincidence properties (κ_W , and CV_W) did not vary much with size. The temporal precision of the population oscillation, however, increases approximately as $1/\sqrt{N}$ (Fig. 10h). Large networks can thus produce precise pacemaker rhythms. In addition the statistical quantities in small networks show more variation with different realizations of the current drive. It is of considerable interest to understand why weak synchronization is so much more robust and prevalent compared to strong synchronization. In strong synchronization one requires an equal firing rate for each neuron, and coincident spikes, while weak synchronization requires only coincident spikes. By definition, then, weak synchronization is easier to generate. In fact strong synchronization is intrinsically less robust, for it requires phase locking between neuron pairs. This is only possible (depending on intrinsic properties) for a small difference in driving currents. There is a price to pay, there will be a small phase difference between the firings of each neuron. Pairs with a large phase difference are less stable against the influence of noise. It is

instructive to consider this problem as the nonlinear dynamics of a single neuron driven by a periodic drive. The neuron can be entrained on different $n:m$ steps. On these steps in the $f-I$ plot the neuron generates n action potentials during m cycles of the periodic drive for a range of values of I (or intrinsic frequency). Noise induces jitter in the spike time, but the CV on the steps is reduced compared to the CV outside the steps (and steps with higher m values). The entrainment produces a phase difference between the firing time and the crest of the drive. The size of the phase difference depends on the intrinsic frequency of the neuron. Robustness is less for current values close to the edge of the step (with unfavorable phase differences). Therefore noise will reduce the width of the phase-locking step (not shown). Here we found network parameters for which coincidence could be maintained despite highly variable firing rates of the neuronal populations.

Our work, and also a recent study [21], is to a large extent based on the recent contributions by Wang & Buzsaki [20]. It is therefore important to briefly reiterate, and spell out how our work extends the work of Wang & Buzsaki, and how it differs [21]. We included the effect of synaptic noise. We have shown that for the purposes of our modeling work a Gaussian white noise current can adequately reproduce experimental ranges of CV [46]. Biophysically realistic amounts of noise do affect the synchronization we have studied. The noise effects are also different from those of current heterogeneity as was discussed before. The issue of noise was not addressed by Wang & Buzsaki. Another important difference is that previous works [20,21] studied only strong synchronization. Here we have proposed that stochastic weak synchronization underlies the synchronized population oscillations in the hippocampus. For this reason our robust 40Hz population rhythms were obtained for different values of the coupling parameters g_{syn} , τ_{syn} , and the driving current I , and the system size N compared to previous work [20]. In our computational work we actually needed a small amount of noise to obtain weak synchronization.

The synchronization properties of large networks may be of some mathematical interest. Our networks are small, and probably the behavior can change quantitatively when increasing the network size significantly. However, in this paper we only addressed the question as to whether networks of physiologically realistic size and connectivity can robustly synchronize. Recent experimental work suggests that interneurons contact on the order of 60 other interneurons [54]. For that reason we only vary our network size between $N = 10$ and 1000.

In the introduction we mentioned recent *in vivo* work and the controversies on the functional role of synchronization. Our work obviously does not contribute to the understanding of the function of synchronization. An important question is what kind of synchronization can be sustained in biophysically realistic networks. Traub and coworkers [12,15,14,13] looked for physiological correlates of the gamma rhythms using *in vitro* and computational experiments. Their results show the crucial role of inhibition, and have provided much of the impetus for our work. The nervous system produces, for some unknown reason, periodic population activity using

circuitry consisting of noisy and heterogeneous neurons. Our results establish that it is possible for inhibitory neurons to be the driving force for synchronization under these conditions.

V. ACKNOWLEDGMENTS

This work was partially funded by the Northeastern University CIRCS fund, and the Sloan Center for Theoretical Neurobiology (PT). We thank W-J Rappel for help during the initial stage of this work, and TJ Sejnowski for useful suggestions. Part of the calculations were performed at Northeastern University High Performance Computer Center.

-
- [1] Adrian ED. The electrical activity of the mammalian olfactory bulb. *Electroencephalogr. Clin. Neurophysiol.*, 2:377–388, 1950.
 - [2] Softky WR and Koch C. The highly irregular firing of cortical cells is inconsistent with temporal integration of random EPSPs. *J. Neurosci.*, 13:334–350, 1993.
 - [3] Gur M Beylin A and Snodderly DM. Response variability of neurons in primary visual cortex (V1) of alert monkeys. *J. Neurosci.*, 17:2914–2920, 1997.
 - [4] Aertsen A and Arndt M. Response synchronization in the visual cortex. *Curr. Opin. Neurobiol.*, 3:586–594, 1993.
 - [5] Rieke F, Warland D, de Ruyter van Steveninck RR, and Bialek W. *Spikes: exploring the neural code*. MIT press, Cambridge, 1997.
 - [6] Softky WR. Simple codes versus efficient codes. *Curr. Opin. Neurobiol.*, 5:239–247, 1995.
 - [7] Shadlen MN and Newsome WT. Noise, neural codes, and cortical organization. *Curr. Opin. Neurobiol.*, 4:569–579, 1994.
 - [8] Shadlen MN and Newsome WT. Is there signal in the noise (comment)? *Curr. Opin. Neurobiol.*, 5:248–250, 1995.
 - [9] Shadlen MN and Newsome WT. The variable discharge of cortical neurons: implications for connectivity, computation, and information coding. *J. Neurosci.*, 18:3870–3896, 1998.
 - [10] Stopfer M, Bhagavan S, Smith BH, and Laurent G. Impaired odor discrimination on desynchronization of odor-encoding neural assemblies. *Nature*, 390:70–74, 1997.
 - [11] Prut Y, Vaadia E, Bergman H, Haalman I, Slovin H, and Abeles M. Spatiotemporal structure of cortical activity: properties and behavioral relevance. *J. Neurophysiol.*, 79:2857–2874, 1998.
 - [12] Traub RD, Whittington MA, Colling SB, Buzsaki G, and Jeffreys JGR. Analysis of gamma rhythms in the rat hippocampus in vitro and in vivo. *J. Physiol.*, 493:471–484, 1996.
 - [13] Whittington MA, Traub RD, Faulkner HJ, Jefferys JGR, and Chettiar K. Morphine disrupts long-range synchrony of gamma oscillations in hippocampal slices. *Proc. Natl. Acad. Sci.*, 95:5807–5811, 1998.
 - [14] Whittington MA, Traub RD, and Jeffreys JGR. Synchronized oscillations in interneuron networks driven by metabotropic glutamate receptor activation. *Nature*, 373:612–615, 1995.
 - [15] Traub RD, Whittington MA, Stanford IM, and Jeffreys JGR. A mechanism for generation of long-range synchronous fast oscillations in the cortex. *Nature*, 383:621–624, 1996.
 - [16] Fisahn A, Pike FG, Buhl EH, and Paulsen O. Cholinergic induction of network oscillations at 40 Hz in the hippocampus in vitro. *Nature*, 394:186–189, 1998.
 - [17] Fellous J-M and Sejnowski TJ. The involvement of CA1 and CA3 in Carbachol-induced EEG oscillations in the hippocampal slice. *Society of Neuroscience Abstract*, 1998.
 - [18] van Vreeswijk C, Abbott LF, and Ermentrout GB. When inhibition not excitation synchronizes neural firing. *J. Comput. Neurosci.*, 1:313–322, 1994.
 - [19] Wang X-J. Ionic basis for intrinsic 40 Hz neuronal oscillations. *Neuroreport*, 5:221–224, 1993.
 - [20] Wang XJ and Buzsáki G. Gamma oscillation by synaptic inhibition in a hippocampal interneuronal network model. *J. Neurosci.*, 16:6402–6413, 1996.
 - [21] White JA, Chow CC, Ritt J, Soto-Treviño C, and Kopell N. Synchronization and oscillatory dynamics in heterogeneous, mutually inhibited neurons. *J. Comput. Neurosci.*, 5:5–16, 1998.
 - [22] Tiesinga PHE, Rappel W-J, and José, JV. Synchronization in networks of noisy interneurons. In Bower J, editor, *Computational Neuroscience*, pages 555–559. Plenum Press, New York, 1998.
 - [23] Brunel N and Hakim V. Fast global oscillations in networks of integrate-and-fire neurons with low firing rates. *preprint*, 1998.
 - [24] Amit DJ and Brunel N. Model of global spontaneous activity and local structured activity during delay periods in the cerebral cortex. *Cereb. Cortex*, 7:237–252, 1997.
 - [25] Golomb D and Rinzel J. Clustering in globally coupled inhibitory neurons. *Physica D*, 72:259–282, 1994.
 - [26] Golomb D and Rinzel J. Dynamics of globally coupled inhibitory neurons with heterogeneity. *Phys. Rev. E*, 48:4810–4814, 1993.
 - [27] Golomb D and Rinzel J. Synchronization properties of spindle oscillations in a thalamic reticular nucleus model. *J. Neurophys.*, 72:1109–1126, 1994.
 - [28] Kopell N and LeMasson G. Rhythmogenesis, amplitude modulation, and multiplexing in a cortical architecture. *Proc. Natl. Acad. Sci.*, 91:10586–10590, 1994.
 - [29] Rall W. Perspective on Neuron Model Complexity. In Arbib, MA, editor, *Handbook of Brain theory and neural networks*, pages 728–732. MIT press, 1995.
 - [30] Pinsky PF and Rinzel J. Intrinsic and network rhythmogenesis in a reduced Traub model for CA3 neurons. *J. Comput. Neurosci.*, 1:39–60, 1994.
 - [31] Mainen ZF and Sejnowski TJ. Influence of dendritic structure on firing pattern in model neocortical neurons. *Nature*, 383:363–366, 1994.
 - [32] Mainen ZF and Sejnowski TJ. Modeling active dendritic processes in pyramidal neurons. In Koch C and Segev I, editors, *Methods in neuronal modeling*, pages 171–209. MIT press, Cambridge, 1998.
 - [33] Yuste R and Tank DW. Dendritic integration in mammalian neurons, a century after Cajal. *Neuron*, 16:701–716, 1996.
 - [34] Koch C. Computation and the single neuron. *Nature*, 385:207–210, 1997.
 - [35] Larkum ME, Zhu JJ, and Sakmann B. A new cellular mechanism for coupling inputs arriving at different cortical papers.

Nature, 398:338–341, 1999.

- [36] Perkel DH, Mulloney B, and Rudelli RW. Quantitative methods for predicting neuronal behavior. *Neuroscience*, 56:823–827, 1981.
- [37] Wang X-J and Rinzel J. Spindle rhythmicity in the reticularis thalami nucleus: synchronization among mutually inhibitory neurons. *Neuroscience*, 53:899–904, 1993.
- [38] Buhl EH, Cobb SR, Halasy K, and Somogyi P. Properties of unitary IPSPs evoked by anatomically identified basket cells in the rat hippocampus. *Eur. J. Neurosci.*, 7:1989–2004, 1995.
- [39] Greenside HS and Helfand E. Numerical integration of stochastic differential equations. *Bell Syst. Tech. J.*, 60:1927, 1981.
- [40] Press WH, Teukolsky SA, Vetterling WT, and Flannery BP. *Numerical Recipes*. Cambridge University Press, Cambridge, 1992.
- [41] Lacaille J-C, Mueller AL, Kunkel DD, and Schwartzkroin PA. Local circuit interactions between oriens-alveus interneurons and CA1 pyramidal cells in hippocampal slices: electrophysiology and morphology. *J. Neurosci.*, 7:1979–1993, 1987.
- [42] Lacaille J-C and Williams S. Membrane properties of interneurons in stratum oriens-alveus of the CA1 region of rat hippocampus in vitro. *Neuroscience*, 36:349–359, 1990.
- [43] Morin F, Beaulieu, and Lacaille J-C. Membrane properties and synaptic currents evoked in CA1 interneuron subtypes in rat hippocampal slices. *J. Neurophys.*, 76:1–16, 1996.
- [44] Johnston D and Wu M. *Foundations of cellular neurophysiology*. MIT Press, Cambridge, 1995.
- [45] Calvin WH and Stevens CF. Synaptic noise as a source of variability in the the interval between action potentials. *Science*, 155:842–843, 1967.
- [46] Tiesinga PHE and José, JV. Spiking Statistics in Noisy Hippocampal Interneurons. *Proceedings Computational Neuroscience 1998*, 1999.
- [47] Rodieck RW, Kiang NY-S, and Gerstein GL. Some quantitative methods for the study of spontaneous activity of single neurons. *Biophys. J.*, 2:351–368, 1962.
- [48] Gray CM and McCormick DA. Chattering cells: superficial pyramidal neurons contributing to the generation of synchronous oscillations in the visual cortex. *Science*, 274:109–113, 1996.
- [49] Hunter JD, Milton JG, Thomas PJ, and Cowan JD. Resonance effect for neural spike time reliability. *J. Neurophysiol.*, 80:1427–1438, 1998.
- [50] Wiesenfeld K and Moss F. Stochastic resonance and the benefits of noise: from ice ages to crayfish and SQUIDS. *Nature*, 373:33–36, 1995.
- [51] Moortgat KT, Keller CH, Bullock TH, and Sejnowski TJ. Sub-microsecond pacemaker precision is behaviorally modulated: the gymnotiform electromotor pathway. *Proc. Natl. Acad. Sci.*, 95:4684–4689, 1998.
- [52] MacLeod K and Laurent G. Distinct mechanisms for synchronization and temporal patterning of odor-encoding neural assemblies. *Science*, 274:976–979, 1996.
- [53] Singer W and Gray CM. Visual feature integration and the temporal correlation hypothesis. *Annu. Rev. Neurosci.*, pages 555–586, 1995.
- [54] Sik A, Penttonen M, Ylinen A, and Buzsáki G. Hippocampal CA1 interneurons: An in vivo intracellular labeling study. *J. Neurosci.*, 15:6651–6665, 1995.




FIG. 1. Coherence parameter κ (dashed line, left hand scale), and average network frequency f (continuous line, right hand scale) versus (a) noise strength D (with $\sigma_I = 0$) and (b) current heterogeneity σ_I (with $D = 0$). After a transient of 1 s, (a) time-averages were computed over 3 s, and (b) 2 s. In (b) each point represents the results of one independent value of the current heterogeneity. In (c,d) we plot the synaptic drive $s(t)$, (c) for $\sigma_I = 0$ with $D = 0, 0.01, 0.04$, and 0.09 from top to bottom; whereas in (d) $D = 0$ with $\sigma_I = 0, 0.01, 0.04$, and 0.09 from top to bottom. The curves are offset by multiples of $\Delta s = 1$. We used the standard set of parameters described in Methods.




FIG. 2. (I) Loss of synchrony due to current heterogeneities: (a) $\sigma_I = 0.02$, (b) 0.07 , (c) 0.08 , respectively. (II) Loss of synchrony due to synaptic noise: $D = 0.01$ (a), 0.04 (b), 0.09 (c), respectively. In 1 and 13 we plot the autocorrelation of X_1 (spike train of neuron 1); In 2 – 10 we plot the cross correlation of X_1 with $X_2, \dots, X_{10}$, respectively; In 11 we show the cross correlations between X_i and X_j , averaged over all pairs i, j ; In 12 we plot the autocorrelation of X_i , averaged over all neurons, and in 14 the auto correlation of the total spiking rate X ; In 15 we show the rastergrams of the network, i.e. neuron number versus spiking time. The time-scale bar, shown in Ic12, applies to the curves 1 – 14, (a-c) in I and II. The y-axis is in arbitrary units, and the same scale is used for the curves in 1 – 11, and 12 – 14, except for the curves IIa 12,13 and IIb 12,13, which are rescaled by a factor of 10. After a transient of 1 s, the time averages are computed over 2 s (I), and 3 s (II). We used the standard set of parameters described in Methods.




FIG. 3. (a-f) distribution of κ values for individual pairs (i,j) , (g-i) difference in driving current $\Delta I = |I_i - I_j|$ versus κ . In (a-c) $\sigma_I = 0$ and (a) $D = 0.01$, (b) 0.04 , (c) 0.09 . Whereas in (d-i) $D = 0$ and (d,g) $\sigma_I = 0.02$, (e,h) 0.04 , and (f,i) 0.15 . A transient of 500 ms was discarded before averaging over 2000 ms. We used the standard set of parameters described in Methods.




FIG. 4. (a) Average interspike interval τ_{ISI} , and (b) coefficient of variation (CV), plotted versus noise strength D . We consider four different cases of ISIHs, labeled by numbers in (c)-(e) and by symbols in (a) and (b), (1) (circles) averages over all neurons in the network, (2) one neuron in the network, (3) (diamonds) isolated neuron with autosynaptic feedback, and (4) (squares) isolated neuron without feedback. The noise strengths are (c) $D = 0.01$, (d) 0.04 , and (e) 0.09 , respectively. In all graphs $I = 1.0$ except in case (4), with $I = 0.61$. The time-scale bar for the ISIH is shown in c 1, the y-scale is arbitrary, but it is the same for all curves with the same D -value. The averaging times are 10 s (network), and 200 s (isolated neuron).




FIG. 5. (a) Coherence parameter κ , (b) the coefficient of variation CV for a network, and (c) CV for an isolated neuron, versus the applied current I for different values of $D = 0.0$ (continuous line), 0.004 (dotted line), 0.04 (dashed line), and 0.2 (dot-dashed line). ISI histograms for a network of neurons (1,3), and an isolated neuron (2,4) for different values of the current (d) $I = 0.8$, (e) 0.96 , and (f) 1.14 . We used $D = 0.0$ (1,2) and 0.004 (3,4). The bar time-scale for the ISI histogram is shown in d2. We applied an excitatory drive with reversal potential $E = 0 \text{ mV}$, strength $g_{syn} = 0.05$, and frequency 40 Hz . The isolated neuron has an autosynaptic feedback of $g_{syn} = 0.1$. After a transient of 1 s (0.5 s), time-averages were calculated over 10 s (50 s) for the network (isolated neuron).




FIG. 6. (a) Network frequency, f , and (b) synchronization parameter, κ , plotted versus noise strength D , for different values of the synaptic coupling and applied current $(g_{syn}, I) = (0.02, 0.6955)$, $(0.1, 1.0)$, $(0.3, 1.625)$, and $(0.5, 2.15)$, labeled by (1)-(4), respectively. The value of the applied current is chosen such that the neuron network will fire at $f \approx 39.05$ Hz at $D = 0$. In (c) we plot κ versus g_{syn} for $D = 0.02$ at the aforementioned current values. Time averages are computed over 10 s after a transient of 1 s,




FIG. 7. Rastergrams comparison, (a) $g_{syn} = 0.1$, $I = 1.0$, and (b) $g_{syn} = 0.5$, $I = 2.15$, with noise strength $D = 0.002$. We also compare the ISI histograms for different values of the synaptic coupling $g_{syn} = 0.5$ (1), 0.2 (2), and 0.05 (3), and for different values of the noise strength D equal to (c) 0.014 , (d) 0.034 , (e) 0.07 (e), and (f) 0.17 . The values of the applied currents are $I = 2.15$, 1.331 , and 0.8145 for (1), (2), and (3), respectively. Time-scale bar is shown in f1, the y-scale is arbitrary but the same for all curves in (c)-(f). For clarity the top of c1 and d1 is cut off. An initial transient of 10 s in (a) and (b) was discarded. After a transient of 1 s, time-averages were taken over 10 s (c-f).




FIG. 8. We plot (a) network frequency f_n , (b) single neuron firing rate f , (c) cluster size N_c normalized by system size $N = 100$, (d) κ , (e) coefficient of variation CV of ISI, and (f) standard deviation of the network period τ_n versus synaptic strength g_{syn} , for different values of $D = 0.0$ (dot-dashed line), $D = 0.008$ (dotted) and $D = 0.2$ (solid line). g_{syn} is varied from 0.05 to 2.5 in increments of 0.05. For $D = 0.008$ ($D = 0.2$) the first 6 (14) points have been removed according to the criterium given in the Methods section. In (b) we have added the firing rate of a single neuron with autosynaptic feedback (fat solid line), and the network for $D = 0$ (open circles). Other parameters are $\sigma_I = 0$, $I = 2.0$, $\tau_{syn} = 20$. A 500 ms transient is discarded, and averages are taken over 2000 ms. To smooth (d-f) we have performed running averages over five points, the original points are denoted by dots.




FIG. 9. We plot (a)-(e) as in the previous figure, (f) number N_s of active neurons for different values of $D = 0.0$ (solid), $D = 0.008$ (dot-dashed). Other parameters are $\sigma_I = 0.02$, $I = 2.0$, $\tau_{syn} = 20$. The smoothed (d-f) figures were obtained as in the previous figure.




FIG. 10. We plot (a) network frequency f_n , (b) single neuron firing rate f , (c) coefficient of variation CV of ISI, (d) κ , (e) cluster size N_c normalized by system size N , (f) number of active neurons N_s normalized by N , (g) coefficient of variation of N_c , (h) standard deviation of the network period τ_n versus system size N ($N = 10, 20, 50, 100, 200, 500, 1000$). We show results for ten different realizations of the current distribution (open circles), and the their average (solid lines). Other parameters are $\sigma_I = 0.10$, $D = 0.2$, $I = 5.0$, $g_{syn} = 1.0$, and $\tau_{syn} = 20$. A 500 ms transient is discarded, and averages are over at least 2000 ms.




FIG. 11. We plot in (a,c) the interspike intervals (ISI) of the network neurons sorted from the lowest to highest value, and (b,d) the ISIh. In (a,c) we used different values of the noise-strength, from top to bottom $D = 0.0, 0.04, 0.08, 0.36$, and 0.56 , with $\sigma_I = 0$. In (c,d) from top to bottom $\sigma_I = 0$ (1), 0.173 (2), 0.346 (3), 0.520 (4), and 0.693 (5), with $D = 0.2$. After a transient of 500 ms, the time-average is computed over 2 s. We plot the instantaneous firing rate as a function of time for (e) $D = 0, \sigma_I = 0$, (f) $D = 0.56, \sigma_I = 0$, and (g) $D = 0.2, \sigma_I = 0.35$. The scale bars are (g) 500 imp/s, (h,i) 50 imp/s. We used $I = 3.5, \tau_{syn} = 20, N = 1000$, and $g_{syn} = 2.0$.




FIG. 12. We plot in (a,d) the network frequency f_n (solid line, left hand scale) and single neuron firing rate f (dashed line, right hand scale); (b,e) coherence κ (solid line) and the CV (dashed line); (c,f) cluster size N_c and number of active neurons N_s divided by N ; as function of (a,b,c) noise variance D , and (d,e,f) current heterogeneity σ_I . We used the following parameters $I = 3.5$, $\tau_{syn} = 20$, $g_{syn} = 0$, and $N = 1000$, for (a,b,c) with $\sigma_I = 0$ and for (d,e,f) the same parameters with $D = 0.2$.




FIG. 13. We plot in (a,b) the interspike intervals (ISI) for the 100 neurons sorted from the lowest to highest values, and in (c,d) the ISI histograms. In (a,c) we used different values for the noise-strength, from top to bottom $D = 0.04, 0.24, 0.84, 4.2$, and 8.2 , with $\sigma_I = 0$. In (b,d) from top to bottom $\sigma_I = 0$ (1), 0.04 (2), 0.08 (3), 0.12 (4), and 0.16 (5), with $D = 1.04$. A sinusoidal current with a period of 25 ms and an amplitude $1.2 \mu A/cm^2$ is added to the soma. The time-scale bar is shown in d1, the y-scale is arbitrary but the same in (c), and (d). After a transient of 2 s, the time-average is computed over 10 s. (e,f) Instantaneous spiking rate versus time for (e) $\sigma_I = 0$, and (f) $\sigma_I = 0.10$, both at $D = 1.04$.